








Amendments to the Claims:

1. (Previously Presented) A compound corresponding to the formula (I):



in which:

X represents a group selected from: glucose, fructose, mannose, galactose, ribose, maltose, glucosamine, sucrose and lactobionamide, a poly(ethylene oxide) chain consisting of from 30 to 100 ethylene oxide units, a group selected from,



m represents an integer equal to 1, 2 or 3;

Y represents a spacer arm which is intended to link the aromatic nucleus to the

hydrophilic X substituents; and

Y is selected from  $\text{—O—}\overset{\text{O}}{\parallel}\text{C—}$ ,  $\text{—NH—}\overset{\text{O}}{\parallel}\text{C—}$ ,  $\text{—NH—}\overset{\text{O}}{\parallel}\text{C—NH—}$ ,  $\text{—O—}\overset{\text{O}}{\parallel}\text{C—NH—}$ ,  $\text{—O—}$ ,  $\text{—S—}$ ,  $\text{—NH—}$ , and  $\text{C}_1\text{—C}_6$  hydrocarbon chains which are optionally interrupted by one or more of the

following groups:  $\text{—O—}\overset{\text{O}}{\parallel}\text{C—}$ ,  $\text{—NH—}\overset{\text{O}}{\parallel}\text{C—}$ ,  $\text{—NH—}\overset{\text{O}}{\parallel}\text{C—NH—}$ ,  $\text{—O—}\overset{\text{O}}{\parallel}\text{C—NH—}$ ,  $\text{—O—}$ ,  $\text{—S—}$ ,  $\text{—NH—}$ ;

y represents an integer equal to 0 or to 1;

Y' represents a group selected from  $\text{—O—}\overset{\text{O}}{\parallel}\text{C—}$ ,  $\text{—NH—}\overset{\text{O}}{\parallel}\text{C—}$ ,  $\text{—NH—}\overset{\text{O}}{\parallel}\text{C—NH—}$ ,  $\text{—O—}\overset{\text{O}}{\parallel}\text{C—NH—}$ ,  $\text{—O—}\overset{\text{O}}{\parallel}\text{C—NH—}$ ,  $\text{—O—}$ ,  $\text{—S—}$ ;


m' is an integer selected from 1 and 2;

X' represents a hydrogen atom or a  $\text{C}_4\text{—C}_{14}$  alkyl chain which is optionally substituted by one or more fluorine atoms.

2. (Previously Presented) The compound as claimed in claim 1, wherein X represents a group selected from: glucose, lactose, manose, galactose, ribose, maltose, glucosamine, sucrose and lactobionamide.

3. (Previously Presented) A compound as claimed in claim 1, wherein X represents a group selected from poly(ethylene oxide) chains consisting of from 50 to 60 units.

4. (Previously Presented) A compound as claimed in claim 1, wherein X represents a group selected from



5. (Previously Presented) A compound as claimed in claim 1, wherein at least one of the following conditions is satisfied:

X represents a group selected from: lacto-bionamide,


m represents 1;

m' represents 1 or 2;

X' is selected from the groups octyl, decyl, dodecyl and  $\text{CF}_3(\text{CF}_2)_r\text{CH}_2\text{CH}_2-$ , where  $8 \geq r \geq 6$ .


6. (Previously Presented) A process for preparing a compound corresponding to the formula (I) as claimed in Claim 1 wherein an aldehyde corresponding to the formula (II) is reacted with a hydroxylamine corresponding to the formula (III) in accordance with scheme 2

below:



Scheme 2

7. (Previously Presented) The process as claimed in claim 6, wherein the compound of the formula (III) is prepared in accordance with a process which is described in scheme 3:



Scheme 3

8. (Previously Presented) A pharmaceutical composition comprising at least one compound corresponding to the formula (I) as claimed in Claim 1 in a pharmaceutically acceptable excipient.

9 – 11 (Cancelled)

12. (Previously Presented) A cosmetic composition, comprising at least one compound corresponding to the formula (I) as claimed in Claim 1 in a cosmetically acceptable

excipient.

13. (Cancelled)

14. (Previously Presented) A method of capturing free radicals comprising the step of reacting a free radical with the compound as claimed in Claim 1.

15. (Previously Presented) A compound as claimed in claim 1, wherein X represents a group selected from: glucosamine, sucrose and lactobionamide.

16. (Previously Presented) The compound as claimed in claim 1, wherein Y represents a group selected from:

